

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/00, 38/04		A1	(11) International Publication Number: WO 99/33482
			(43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/US98/27367 (22) International Filing Date: 23 December 1998 (23.12.98) (30) Priority Data: 60/070,153 31 December 1997 (31.12.97) US 60/080,588 3 April 1998 (03.04.98) US (71) Applicant: UNIVERSITY OF UTAH RESEARCH FOUNDATION [US/US]; Suite 170, 421 Wakara Way, Salt Lake City, UT 84108 (US). (72) Inventors: OLIVERA, Baldomero, M.; 1370 Bryan Avenue, Salt Lake City, UT 84108 (US); McINTOSH, J., Michael; 1151 South 2000 East, Salt Lake City, UT 84108 (US); YOSHIKAMI, Doju; 1074 East 6 South, Salt Lake City, UT 84102 (US); CARTIER, G., Edward; 825 West River Terrace #617, Salt Lake City, UT 84123 (US); LUO, Siqin; 1179 South Blair Street #21, Salt Lake City, UT 84111 (US). (74) Agents: IHNEN, Jeffrey, L. et al.; Rothwell, Figg, Ernst & Kurz, Suite 701 East, 555 13th Street, N.W., Columbia Square, Washington, DC 20004 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report: Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: USES OF ALPHA-CONOTOXIN PEPTIDES			
(57) Abstract <p>the present invention relates to the use of α-conotoxin peptides having the general formula: Xaa₁-Xaa₂-Cys-Cys-Xaa₃-Xaa₄-Pro-Xaa₅-Cys-Xaa₆-Cys (SEQ ID NO.1) for treating disorders regulated at neuronal nicotinic acetylcholine receptors. Such disorders include, but are not limited to, cardiovascular disorders, gastric motility disorders, urinary incontinence, nicotine addiction, mood disorders (such as bipolar disorder, unipolar depression, dysthymia and seasonal effective disorder) and small cell lung carcinoma, as well as the localization of small cell lung carcinoma. In this formula, Xaa₁ is des-Xaa₁, Tyr, mono-iodo-Tyr or di-iodo-Tyr, Xaa₂ is any amino acid, Xaa₃ is any amino acid, Xaa₄ is any amino acid, Xaa₅ is any amino acid and Xaa₆ represents a peptide of 3-7 amino acids. Disulfide linkages exist between the first and third cysteines and the second and fourth cysteines. Pro may be replaced with hydroxy-Pro. The C-terminus may contain a hydroxyl or an amide group, preferably an amide group.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

USES OF ALPHA-CONOTOXIN PEPTIDES

This invention was made with Government support under Grant Nos. GM48677 and
5 MH53631 awarded by the National Institutes of Health, Bethesda, Maryland. The United States
Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

This invention relates to uses of relatively short peptides about 14-17 residues in length,
10 which are naturally available in minute amounts in the venom of the cone snails or analogs to the
naturally available peptides, and which include two cyclizing disulfide linkages.

The publications and other materials used herein to illuminate the background of the
invention, and in particular, cases to provide additional details respecting the practice, are
incorporated by reference, and for convenience are referenced in the following text by author and
15 date and are listed alphabetically by author in the appended bibliography.

Mollusks of the genus *Conus* produce a venom that enables them to carry out their unique
predatory lifestyle. Prey are immobilized by the venom that is injected by means of a highly
specialized venom apparatus, a disposable hollow tooth that functions both in the manner of a
harpoon and a hypodermic needle.

20 Few interactions between organisms are more striking than those between a venomous
animal and its envenomated victim. Venom may be used as a primary weapon to capture prey or
as a defense mechanism. Many of these venoms contain molecules directed to receptors and ion
channels of neuromuscular systems.

The predatory cone snails (*Conus*) have developed a unique biological strategy. Their
25 venom contains relatively small peptides that are targeted to various neuromuscular receptors and
may be equivalent in their pharmacological diversity to the alkaloids of plants or secondary
metabolites of microorganisms. Many of these peptides are among the smallest nucleic acid-
encoded translation products having defined conformations, and as such, they are somewhat
unusual. Peptides in this size range normally equilibrate among many conformations. Proteins
30 having a fixed conformation are generally much larger.

The cone snails that produce these toxic peptides, which are generally referred to as
conotoxins or conotoxin peptides, are a large genus of venomous gastropods comprising
approximately 500 species. All cone snail species are predators that inject venom to capture prey,

and the spectrum of animals that the genus as a whole can envenomate is broad. A wide variety of hunting strategies are used, however, every *Conus* species uses fundamentally the same basic pattern of envenomation.

Several peptides isolated from *Conus* venoms have been characterized. These include the
5 α -, μ - and ω -conotoxins which target nicotinic acetylcholine receptors, muscle sodium channels, and neuronal calcium channels, respectively (Olivera et al., 1985). Conopressins, which are vasopressin analogs, have also been identified (Cruz et al., 1987). In addition, peptides named conantokins have been isolated from *Conus geographus* and *Conus tulipa* (Mena et al., 1990; Haack et al., 1990). These peptides have unusual age-dependent physiological effects: they induce a sleep-
10 like state in mice younger than two weeks and hyperactive behavior in mice older than 3 weeks (Haack et al., 1990). Recently, peptides named contryphans containing D-tryptophan residues have been isolated from *Conus radiatus* (U.S. Serial No. 60/030,722), and bromo-tryptophan conopeptides have been isolated from *Conus imperialis* and *Conus radiatus* (U.S. Serial No. 08/785,534).

15 Neuronal nicotinic acetylcholine receptors (nAChRs) are believed to be heteropentameric ion channel complexes generally requiring at least two different subunits (an α and a β). Molecular data indicate that in the mammalian central nervous system there exists a large number of different nAChR subunits. To date, seven different α subunits ($\alpha 2$ - $\alpha 7$, $\alpha 9$) and three different β subunits ($\beta 2$ - $\beta 4$) have been defined by cloning. The $\alpha 3\beta 4$ -containing nAChR subtype and the $\alpha 3\beta 2$ -containing
20 nAChR subtype are each present in the autonomic nervous system and in the central nervous system. The $\alpha 7$ -containing nAChR subtype is also present in the autonomic nervous system.

While postsynaptic nAChRs have been recognized for some time, more recent data have demonstrated the presence of presynaptic neuronal nAChRs. Agonist stimulation of presynaptic nAChRs induces neurotransmitter release. Nicotinic agonists have been shown to elicit the release
25 of several different neurotransmitters, including dopamine from striatum and frontal cortex (El-Bizri and Clarke, 1994; Grady et al., 1992; Rapier et al., 1988); norepinephrine from hippocampus (Clarke and Reuben, 1996; Rowell and Winkler, 1984; Sacaan et al., 1995; Wilkie et al., 1993); glutamate from cortex, medial habenula nucleus and hippocampus (McGehee and Role, 1995; Vidal and Changeux, 1993; Gray et al., 1996); GABA from interpeduncular nucleus (Mulle et al., 1991)
30 and acetylcholine for cortex and hippocampus (Lapchak et al., 1989; Rowell and Winkler, 1984).

In addition, it appears that distinct subtypes of presynaptic nAChRs regulate the release of different neurotransmitters. For example, nicotine-stimulated glutamate and acetylcholine release

-3-


are blocked by α -bungarotoxin suggesting that these nAChRs include an $\alpha 7$ subunit (McGehee and Role, 1995). In contrast, nicotine-stimulated dopamine release is not blocked by α -bungarotoxin (Grady et al., 1992). Furthermore, the nAChRs modulating norepinephrine release pharmacologically differ from those modulating the release of glutamate, acetylcholine or dopamine (Clarke and Reuben, 1996; Sacaan et al., 1995).

As previously described, presynaptic nAChRs in the central nervous system (CNS) modulate the release of several neurotransmitters, including norepinephrine and dopamine (Wonnacott, 1997). CNS norepinephrine levels are important in the treatment and/or pathophysiology of mood disorders (Schatzberg and Nemeroff, 1995; Mongeau et al., 1997). CNS dopamine levels are important in addictive and psychotic disorders (Pontieri et al., 1996; Kahn and Davis, 1995). Thus, the possibility of selectively modulating the presynaptic release of specific neurotransmitters and the possibility of selectively targeting specific nAChRs has significant therapeutic applications. One example of a therapeutic application is tobacco addiction. Studies of nicotine self-administration in animal models suggest that block of nAChRs decreases the reinforcing properties of nicotine. Examples of therapeutic applications resulting from selectively targeting the nAChRs of the autonomic nervous system are the treatment of cardiovascular disorders, gastric mobility disorders and urinary incontinence.

It is desired to identify additional compounds which target different nAChR subtypes as well as the nAChR subtypes of the autonomic nervous system and the central nervous system. It is further desired to identify compounds which are useful as cardiovascular agents, gastric motility agents, urinary incontinence agents, anti-smoking agents, anti-cancer agents, anti-psychotic agents and anti-mood disorder agents.

SUMMARY OF THE INVENTION

This invention relates to uses of relatively short peptides about 14-17 residues in length, which are naturally available in minute amounts in the venom of the cone snails or analogs to the naturally available peptides, and which include two cyclizing disulfide linkages. More specifically, the present invention relates to the use of α -conotoxin peptides having the general formula



for treating disorders regulated at neuronal nicotinic acetylcholine receptors. Such disorders include, but are not limited to, cardiovascular disorders, gastric motility disorders, urinary incontinence, nicotine addiction, mood disorders (such as bipolar disorder, unipolar depression,

-4-

dysthymia and seasonal effective disorder) and small cell lung carcinoma, as well as the localization of small cell lung carcinoma. In this formula, Xaa₁ is des-Xaa₁, Tyr, mono-iodo-Tyr or di-iodo-Tyr, Xaa₂ is any amino acid, Xaa₃ is any amino acid, Xaa₄ is any amino acid, Xaa₅ is any amino acid and Xaa₆ represents a peptide of 3-7 amino acids, with the proviso that when the disorder is small cell lung carcinoma, then the α -conotoxin peptide is not a peptide having an amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:13. Disulfide linkages exist between the first and third cysteines and the second and fourth cysteines. Pro may be replaced with hydroxy-Pro. The C-terminus may contain a hydroxyl or an amide group, preferably an amide group.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows the selectivity of 3 μ M α -conotoxins AuIA, AuIB and AuIC on nAChRs expressed in oocytes.

Figure 2 shows the antagonistic activity of PnIA (Δ), PnIB (\circ), PnIA A10L (\blacksquare) and PnIA N11S (\blacktriangledown) with respect to the α 7 subtype of neuronal nicotinic acetylcholine receptors.


Figures 3A and 3B show the effects of α -conotoxins on nicotine-stimulated release of norepinephrine from rat hippocampal synaptosomes (3A) or dopamine from rat striatal synaptosomes (3B). * $P \leq 0.001$. Data are from 3-10 experiments with 3-6 replicates within each experiment.

SUMMARY OF THE SEQUENCE LISTING

SEQ ID NO:1 is the generic formula for the α -conotoxin peptides useful for the present invention. SEQ ID NO:2 is α -conotoxin peptide MII. SEQ ID NO:3 is α -conotoxin peptide Tyr-MII. SEQ ID NO:4 is α -conotoxin peptide FAT-MII (MII with FAT at residues 9-11 instead of HLE in MII). SEQ ID NO:5 is α -conotoxin peptide AuIA. SEQ ID NO:6 is α -conotoxin peptide Tyr-AuIA. SEQ ID NO:7 is α -conotoxin peptide AuIB. SEQ ID NO:8 is α -conotoxin peptide AuIC. SEQ ID NO:9 is α -conotoxin peptide PnIA. SEQ ID NO:10 is α -conotoxin peptide PnIA A10L (PnIA with L at residue 10 instead of A in PnIA). SEQ ID NO:11 is α -conotoxin peptide PnIA N11S (PnIA with S at residue 11 instead of N in PnIA). SEQ ID NO:12 is α -conotoxin peptide PnIB. SEQ ID NO:13 is α -conotoxin peptide ImI.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

This invention relates to uses of relatively short peptides about 14-17 residues in length, which are naturally available in minute amounts in the venom of the cone snails or analogs to the naturally available peptides, and which include two cyclizing disulfide linkages. More specifically, the present invention relates to the use of α -conotoxin peptides having the general formula



for treating disorders regulated at neuronal nicotinic acetylcholine receptors. Such disorders include, but are not limited to, cardiovascular disorders, gastric motility disorders, urinary incontinence, nicotine addiction, mood disorders (such as bipolar disorder, unipolar depression, dysthymia and seasonal effective disorder) and small cell lung carcinoma, as well as the localization of small cell lung carcinoma. In this formula, Xaa₁ is des-Xaa₁, Tyr, mono-iodo-Tyr or di-iodo-Tyr, Xaa₂ is any amino acid, Xaa₃ is any amino acid, Xaa₄ is any amino acid, Xaa₅ is any amino acid and Xaa₆ represents a peptide of 3-7 amino acids, with the proviso that when the disorder is small cell lung carcinoma, then the α -conotoxin peptide is not a peptide having an amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:13. Disulfide linkages exist between the first and third cysteines and the second and fourth cysteines. Pro may be replaced with hydroxy-Pro. The C-terminus may contain a carboxyl or an amide group, preferably an amide group. The amino acid or the amino acid residues of the peptides is an α -amino acid, which includes natural amino acids, including unusual amino acids such as γ -carboxyglutamic acid, as well as modified or non-natural amino acids, such as those described in, for example, Roberts et al. (1983).

Examples of α -conotoxin peptides falling within the generic formula are set forth in Table 1. These conotoxin peptides are members of the α 4 subclass of α -conotoxin peptides. The tyrosine residue at the N-terminus can also be added to the other peptides shown in Table 1. This tyrosine residue can be iodinated to contain 1 or 2 iodines. In addition, the proline residues can be replaced by hydroxyproline. The tryptophan residue may be replaced by bromo-tryptophan. These changes to the peptides do not change the activity of the native peptide.

-6-

TABLE 1

 α -Conotoxin Peptides

	Peptide	Sequence	SEQ ID NO:
	MII	GCCSNPVCHLEHSNLC	2
5	Tyr-MII	YGCCSNPVCHLEHSNLC	3
	FAT-MII	GCCSNPVCFATHSNLC	4
	AuIA	GCCSYPPCFATNSDYC	5
	Tyr-AuIA	YGCCSYPPCFATNSDYC	6
	AuIB	GCCSYPPCFATNSD-C	7
10	AuIC	GCCSYPPCFATNSGYC	8
	PnIA	GCCSLPPCAANNPDYC	9
	PnIA A10L	GCCSLPPCALNNPDYC	10
	PnIA N11S	GCCSLPPCAASNPDYC	11
	PnIB	GCCSLPPCALSNPDYC	12
15	ImI	GCCSDPRCA---W-RC	13

Additional peptides falling within the general formula can be made based on the peptides shown in Table 1 by making analogs of these peptides or by making conservative substitutions for the amino acid residues shown in Table 1. For example, a FAT-PnIA analog can be made in which FAT replaces AAN at residues 9-11. Conservative substitutions are well known in the art and include, for example, the change of (or vice versa): alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glycine to proline; isoleucine to leucine or valine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; tryptophan to tyrosine. Changes in these peptides can be made in view of the teachings set forth in U.S. provisional patent application numbers 60/062,783 and 60/065,814 and in international patent application number PCT/US98/22368.

The α -conotoxin peptides of the general formula are active on the $\alpha 3\beta 4$, $\alpha 3\beta 2$ or $\alpha 7$ containing subtypes of nAChRs which are present in the autonomic and central nervous systems. For example, the $\alpha 3\beta 4$ -containing subtype is present in sympathetic ganglia and the central nervous system. Similarly, the $\alpha 3\beta 2$ -containing subtype is present in sympathetic ganglia and the central nervous system. Finally, the $\alpha 7$ -containing subtype is present at peripheral ganglia and in the central nervous system. Thus, they are useful as cardiovascular agents, gastric motility agents, urinary incontinence agents, anti-smoking agents and for the treatment or localization of small cell lung carcinoma. These peptides are also useful for treating psychosis, pain and spastic disorders. Finally, these peptides are useful for treating mood disorders.

The α -conotoxin peptides can be designed to be more specific for one of these subtypes of nAChRs. For example, MII has a higher specificity for the $\alpha 3\beta 2$ -containing subtype, whereas FAT-MII has a higher specificity for the $\alpha 3\beta 2$ -containing subtype. Similarly, PnIA has a higher

-7-

specificity for the $\alpha 3\beta 2$ -containing subtype, whereas PnIA A10L has a higher specificity for the $\alpha 7$ -containing subtype. The peptides set forth in Table 1 have the following specificities (with respect to higher specificity, generally by several orders of magnitude): $\alpha 3\beta 2$ -containing subtype: MII, Tyr-MII, PnIA and PnIA N11S; $\alpha 3\beta 4$ -containing subtype: AuIA, AuIB, AuIC, FAT-MII and Tyr-FAT-MII; and, $\alpha 7$ -containing subtype: PnIB, ImI and PnIA A10L. The specificity of each peptide is readily determined by assaying for subtype specificity in accordance with techniques well known in the art.

These peptides are sufficiently small to be chemically synthesized. General chemical syntheses for preparing the foregoing conopeptides peptides are described hereinafter, along with specific chemical synthesis of conopeptides and indications of biological activities of these synthetic products. Various ones of these conopeptides can also be obtained by isolation and purification from specific *Conus* species using the technique described in U.S. Patent No. 4,447,356 (Olivera et al., 1984), the disclosure of which is incorporated herein by reference.

Although the native conopeptides can be obtained by purification from cone snails, because the amounts of conopeptides obtainable from individual snails are very small, the desired substantially pure conopeptides are best practically obtained in commercially valuable amounts by chemical synthesis using solid-phase strategy. For example, the yield from a single cone snail may be about 10 micrograms or less of conopeptide. By "substantially pure" is meant that the peptide is present in the substantial absence of other biological molecules of the same type; it is preferably present in an amount of at least about 85% purity and preferably at least about 95% purity. Chemical synthesis of biologically active conopeptides depends of course upon correct determination of the amino acid sequence. Thus, the conopeptides of the present invention may be isolated, synthesized and/or substantially pure.

The conopeptides can also be produced by recombinant DNA techniques well known in the art. Such techniques are described by Sambrook et al. (1979). Nucleic acid sequences coding for the α -conotoxin peptides and α -conotoxin propeptides can be isolated and cloned using conventional techniques. Alternatively, nucleic acid sequences coding for the α -conotoxin peptides can be synthesized on the basis of the amino acid sequences of the peptides disclosed herein and the known degeneracy of the genetic code. The nucleic acids for the peptides can be designed to achieve maximal expression in a given host system. The peptides produced in this manner are isolated, reduced if necessary, and oxidized to form the correct disulfide bonds, if present in the final molecule.

One method of forming disulfide bonds in the conopeptides of the present invention is the air oxidation of the linear peptides for prolonged periods under cold room temperatures or at room temperature. This procedure results in the creation of a substantial amount of the bioactive, disulfide-linked peptides. The oxidized peptides are fractionated using reverse-phase high performance liquid chromatography (HPLC) or the like, to separate peptides having different linked configurations. Thereafter, either by comparing these fractions with the elution of the native material or by using a simple assay, the particular fraction having the correct linkage for maximum biological potency is easily determined. It is also found that the linear peptide, or the oxidized product having more than one fraction, can sometimes be used for *in vivo* administration because the cross-linking and/or rearrangement which occurs *in vivo* has been found to create the biologically potent conopeptide molecule. However, because of the dilution resulting from the presence of other fractions of less biopotency, a somewhat higher dosage may be required.

The peptides are synthesized by a suitable method, such as by exclusively solid-phase techniques, by partial solid-phase techniques, by fragment condensation or by classical solution couplings.

In conventional solution phase peptide synthesis, the peptide chain can be prepared by a series of coupling reactions in which constituent amino acids are added to the growing peptide chain in the desired sequence. Use of various coupling reagents, e.g., dicyclohexylcarbodiimide or diisopropylcarbonyldimidazole, various active esters, e.g., esters of N-hydroxyphthalimide or N-hydroxy-succinimide, and the various cleavage reagents, to carry out reaction in solution, with subsequent isolation and purification of intermediates, is well known classical peptide methodology. Classical solution synthesis is described in detail in the treatise, "Methoden der Organischen Chemie (Houben-Weyl): Synthese von Peptiden," (1974). Techniques of exclusively solid-phase synthesis are set forth in the textbook, "Solid-Phase Peptide Synthesis," (Stewart and Young, 1969), and are exemplified by the disclosure of U.S. Patent 4,105,603 (Vale et al., 1978). The fragment condensation method of synthesis is exemplified in U.S. Patent 3,972,859 (1976). Other available syntheses are exemplified by U.S. Patents No. 3,842,067 (1974) and 3,862,925 (1975). The synthesis of peptides containing γ -carboxyglutamic acid residues is exemplified by Rivier et al. (1987), Nishiuchi et al. (1993) and Zhou et al. (1996).

Common to such chemical syntheses is the protection of the labile side chain groups of the various amino acid moieties with suitable protecting groups which will prevent a chemical reaction from occurring at that site until the group is ultimately removed. Usually also common is the

protection of an α -amino group on an amino acid or a fragment while that entity reacts at the carboxyl group, followed by the selective removal of the α -amino protecting group to allow subsequent reaction to take place at that location. Accordingly, it is common that, as a step in such a synthesis, an intermediate compound is produced which includes each of the amino acid residues located in its desired sequence in the peptide chain with appropriate side-chain protecting groups linked to various ones of the residues having labile side chains.

As far as the selection of a side chain amino protecting group is concerned, generally one is chosen which is not removed during deprotection of the α -amino groups during the synthesis. However, for some amino acids, e.g., His, protection is not generally necessary. In selecting a particular side chain protecting group to be used in the synthesis of the peptides, the following general rules are followed: (a) the protecting group preferably retains its protecting properties and is not split off under coupling conditions, (b) the protecting group should be stable under the reaction conditions selected for removing the α -amino protecting group at each step of the synthesis, and (c) the side chain protecting group must be removable, upon the completion of the synthesis containing the desired amino acid sequence, under reaction conditions that will not undesirably alter the peptide chain.

It should be possible to prepare many, or even all, of these peptides using recombinant DNA technology. However, when peptides are not so prepared, they are preferably prepared using the Merrifield solid-phase synthesis, although other equivalent chemical syntheses known in the art can also be used as previously mentioned. Solid-phase synthesis is commenced from the C-terminus of the peptide by coupling a protected α -amino acid to a suitable resin. Such a starting material can be prepared by attaching an α -amino-protected amino acid by an ester linkage to a chloromethylated resin or a hydroxymethyl resin, or by an amide bond to a benzhydrylamine (BHA) resin or paramethylbenzhydrylamine (MBHA) resin. Preparation of the hydroxymethyl resin is described by Bodansky et al. (1966). Chloromethylated resins are commercially available from Bio Rad Laboratories (Richmond, CA) and from Lab. Systems, Inc. The preparation of such a resin is described by Stewart and Young (1969). BHA and MBHA resin supports are commercially available, and are generally used when the desired polypeptide being synthesized has an unsubstituted amide at the C-terminus. Thus, solid resin supports may be any of those known in the art, such as one having the formulae $-O-CH_2$ -resin support, $-NH$ BHA resin support, or $-NH$ -MBHA resin support. When the unsubstituted amide is desired, use of a BHA or MBHA resin is preferred,

-10-

because cleavage directly gives the amide. In case the N-methyl amide is desired, it can be generated from an N-methyl BHA resin. Should other substituted amides be desired, the teaching of U.S. Patent No. 4,569,967 (Komreich et al., 1986) can be used, or should still other groups than the free acid be desired at the C-terminus, it may be preferable to synthesize the peptide using classical methods as set forth in the Houben-Weyl text (1974).

The C-terminal amino acid, protected by Boc or Fmoc and by a side-chain protecting group, if appropriate, can be first coupled to a chloromethylated resin according to the procedure set forth in Horiki et al. (1978), using KF in DMF at about 60°C for 24 hours with stirring, when a peptide having free acid at the C-terminus is to be synthesized. Following the coupling of the BOC-protected amino acid to the resin support, the α -amino protecting group is removed, as by using trifluoroacetic acid (TFA) in methylene chloride or TFA alone. The deprotection is carried out at a temperature between about 0°C and room temperature. Other standard cleaving reagents, such as HCl in dioxane, and conditions for removal of specific α -amino protecting groups may be used as described in Schroder and Lubke (1965).

After removal of the α -amino-protecting group, the remaining α -amino- and side chain-protected amino acids are coupled step-wise in the desired order to obtain the intermediate compound defined hereinbefore, or as an alternative to adding each amino acid separately in the synthesis, some may be coupled to one another prior to addition to the solid phase reactor. Selection of an activating reagent is within the skill of the art. Particularly suitable as a coupling reagent is N,N'-dicyclohexylcarbodiimide (DCC, DIC, HBTU, HATU, TBTU in the presence of HoBt or HoAt).

The activating reagents used in the solid phase synthesis of the peptides are well known in the peptide art. Examples of suitable activating reagents are carbodiimides, such as N,N'-diisopropylcarbodiimide and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide. Other activating reagents and their use in peptide coupling are described by Schroder and Lubke (1965) and Kapoor (1970).

Each protected amino acid or amino acid sequence is introduced into the solid-phase reactor in about a twofold or more excess, and the coupling may be carried out in a medium of dimethylformamide (DMF):CH₂Cl₂ (1:1) or in DMF or CH₂Cl₂ alone. In cases where intermediate coupling occurs, the coupling procedure is repeated before removal of the α -amino protecting group prior to the coupling of the next amino acid. The success of the coupling reaction at each stage of the synthesis, if performed manually, is preferably monitored by the ninhydrin reaction, as described

by Kaiser et al. (1970). Coupling reactions can be performed automatically, as on a Beckman 990 automatic synthesizer, using a program such as that reported in Rivier et al. (1978).

After the desired amino acid sequence has been completed, the intermediate peptide can be removed from the resin support by treatment with a reagent, such as liquid hydrogen fluoride or TFA (if using Fmoc chemistry), which not only cleaves the peptide from the resin but also cleaves all remaining side chain protecting groups and also the α -amino protecting group at the N-terminus if it was not previously removed to obtain the peptide in the form of the free acid. If Met is present in the sequence, the Boc protecting group is preferably first removed using trifluoroacetic acid (TFA)/ethanedithiol prior to cleaving the peptide from the resin with HF to eliminate potential S-alkylation. When using hydrogen fluoride or TFA for cleaving, one or more scavengers such as anisole, cresol, dimethyl sulfide and methylethyl sulfide are included in the reaction vessel.

Cyclization of the linear peptide is preferably affected, as opposed to cyclizing the peptide while a part of the peptido-resin, to create bonds between Cys residues. To effect such a disulfide cyclizing linkage, fully protected peptide can be cleaved from a hydroxymethylated resin or a chloromethylated resin support by ammonolysis, as is well known in the art, to yield the fully protected amide intermediate, which is thereafter suitably cyclized and deprotected. Alternatively, deprotection, as well as cleavage of the peptide from the above resins or a benzhydrylamine (BHA) resin or a methylbenzhydrylamine (MBHA), can take place at 0°C with hydrofluoric acid (HF) or TFA, followed by oxidation as described above.

The present α -conotoxins block $\alpha 3\beta 4$ -containing nAChRs, $\alpha 3\beta 2$ -containing nAChRs or $\alpha 7$ -containing nAChRs expressed in *Xenopus* oocytes, as noted above. The present α -conotoxins also block other nAChR subunit combinations but with much lower affinities. For example, at low submicromolar concentrations, e.g., 0.3-1.0 μ M, the AulA blocks essentially only $\alpha 3\beta 4$ -containing receptors (copending application Serial No. 08/857,068, incorporated herein by reference). It is also known that α -conotoxin MII blocks native $\alpha 3\beta 2$ -containing nAChRs and $\alpha 3\beta 4$ -containing nAChRs (copending application Serial No. 08/761,674, incorporated herein by reference).

A particular advantage of α -conotoxin antagonists is their ability to discriminate between nonsymmetrical ligand binding interfaces present on the receptor. The best-studied example is α -conotoxin M1 binding to the muscle nicotinic receptor. In mouse muscle, α -conotoxin M1 displays a four order-of-magnitude selectivity for the $\alpha 1/\delta$ vs. the $\alpha 1/\gamma$ binding site (Sine and Claudio, 1991). Nevertheless, α -conotoxin M1 functionally blocks the muscle receptor with affinity comparable to its affinity for the $\alpha 1/\delta$ binding site, indicating that only one toxin molecule is

-12-

required to prevent channel activation (Martinez et al., 1995). It was also recently demonstrated that α -conotoxin MII has two binding sites on $\alpha 3\beta 2$ and $\alpha 3\beta 4$ containing receptors expressed in *Xenopus* oocytes and only one toxin molecule is required to block function (Cartier et al., 1996b). α -Conotoxin MII discriminates between the $\alpha 3\beta 2$ and $\alpha 3\beta 4$ interface by four orders-of-magnitude (see
5 Serial No. 08/761,674). Thus, α -conotoxin MII has the ability to potently block any receptor containing an $\alpha 3\beta 2$ subunit interface regardless of what other α and β subunits may be present in the receptor complex. α -Conotoxin MII's potency at such receptors would still be high. Similarly, the α -conotoxins AuIA, AuIB and AuIC, have the ability to discriminate between the $\alpha 3\beta 4$ and $\alpha 3\beta 2$ interface (see Serial No. 08/857,068). Consequently, AuIA, AuIB and AuIC have the ability
10 to potently block any receptor containing an $\alpha 3\beta 4$ subunit interface regardless of what other α and β subunits may be present in the receptor complex. These α -conotoxins' potency at such receptors would still be high. Similarly, ImI shows higher specificity to the $\alpha 7$ subunit.

Peptide analogs and peptide mimetics which are specific for the noted subtypes of the nAChR are prepared on the basis of the teachings disclosed herein as well as the teachings presented
15 in the provisional patent application Serial No. 60/065,814 using conventional drug modeling, drug design and combinatorial chemistry. Suitable techniques include, but are not limited to those described in U.S. Patent No. 5,571,698, WO 95/21193, Ecker and Cook (*Bio/Technology* 13:351-360 (1995), Persidis and Persidis (*Bio/Technology* 15:1035-1036 (1997)), Johnson et al. ("Peptide Turn Mimetics" in *Biotechnology and Pharmacy*, Pezzato et al., eds., Chapman and Hall, New York (1993)), Sun and Cohen (*Gene* 137:127-132 (1993)) and the references cited therein. the development of peptide analogs and peptide mimetics are prepared using commercially available
20 drug design software, including those set forth in the Persidis and Persidis reference. These peptide analogs and peptide mimetics have the same activities as the α -conotoxins described herein and in the published literature. As described herein, the specificity of an individual α -conotoxin can be changed by making a peptide analog. The specificity of the peptide analog is determined by using
25 nAChR subtype assays, such as described herein.

Peptide analogs and derivatives can be made in accordance with conventional techniques. Suitable techniques for peptide synthesis is described in U.S. Patent No. 5,514,774, as well as the references cited therein. Peptide mimetics are similarly synthesized by conventional techniques.

30 Pharmaceutical compositions containing a compound of the present invention as the active ingredient can be prepared according to conventional pharmaceutical compounding techniques. See, for example, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton,

PA). Typically, an antagonistic amount of the active ingredient will be admixed with a pharmaceutically acceptable carrier. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., intravenous, oral or parenteral. The compositions may further contain antioxidizing agents, stabilizing agents, preservatives and the like.

5 For oral administration, the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, suspensions or emulsions. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, suspending agents, and the like in the case of oral liquid preparations (such as, for example, 10 suspensions, elixirs and solutions); or carriers such as starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations (such as, for example, powders, capsules and tablets). Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by 15 standard techniques. The active agent can be encapsulated to make it stable to passage through the gastrointestinal tract while at the same time allowing for passage across the blood brain barrier. See for example, WO 96/11698.

For parenteral administration, the compound may be dissolved in a pharmaceutical carrier and administered as either a solution or a suspension. Illustrative of suitable carriers are water, 20 saline, dextrose solutions, fructose solutions, ethanol, or oils of animal, vegetative or synthetic origin. The carrier may also contain other ingredients, for example, preservatives, suspending agents, solubilizing agents, buffers and the like. When the compounds are being administered intrathecally, they may also be dissolved in cerebrospinal fluid.

The active agents, which are peptides, can also be administered in a cell based delivery 25 system in which a DNA sequence encoding an active agent is introduced into cells designed for implantation in the body of the patient, especially in the spinal cord region. Suitable delivery systems are described in U.S. Patent No. 5,550,050 and published PCT Application Nos. WO 92/19195, WO 94/25503, WO 95/01203, WO 95/05452, WO 96/02286, WO 96/02646, WO 96/40871, WO 96/40959 and WO 97/12635. Suitable DNA sequences can be prepared synthetically 30 for each active agent on the basis of the developed sequences and the known genetic code.

The use of α -conotoxin peptides for the treatment or localization of small cell lung carcinoma is disclosed in U.S. Patent No. 5,595,972, incorporated herein by reference. The use of

α -conotoxin peptides for treating cardiovascular disorders is disclosed in international application PCT/US97/20669 designating the U.S., incorporated herein by reference. The use of α -conotoxin peptides for treating nicotine addiction, psychosis and mood disorders is disclosed in U.S. application Serial. No. 08/761,674, incorporated herein by reference. α -Conotoxin peptides with
5 specificity for the $\alpha 3\beta 2$ -containing nAChRs are particularly preferred for treating nicotine addiction. α -Conotoxin peptides with specificity for the $\alpha 3\beta 4$ -containing nAChRs are particularly preferred for treating mood disorders. Gastric motility disorders and urinary incontinence are treated in conventional manner using an antagonistic amount of α -conotoxin peptides disclosed herein.

The α -conotoxin peptides are administered in an amount sufficient to antagonize the $\alpha 3\beta 4$,
10 $\alpha 3\beta 4$ or $\alpha 7$ containing nAChRs as noted above. The dosage range at which the conotoxin peptides exhibit this antagonistic effect can vary widely depending upon the particular condition, e.g., cardiovascular disorders, gastric motility disorders, urinary incontinence, nicotine addiction, mood disorders or small cell lung carcinoma, being treated, the severity of the patient's condition, the patient, the specific conotoxin being administered, the route of administration and the presence of
15 other underlying disease states within the patient. Typically the conopeptides of the present invention exhibit their therapeutic effect at a dosage range from about 0.05 mg/kg to about 250 mg/kg, and preferably from about 0.1 mg/kg to about 100 mg/kg of the active ingredient. A suitable dose can be administered in multiple sub-doses per day. Typically, a dose or sub-dose may contain from about 0.1 mg to about 500 mg of the active ingredient per unit dosage form. A more preferred
20 dosage will contain from about 0.5 mg to about 100 mg of active ingredient per unit dosage form. Dosages are generally initiated at lower levels and increased until desired effects are achieved.

The iodinated analogs of the α -conotoxin peptides can also be used for receptor mapping using conventional techniques. These iodinated analogs can also be used to screen for additional α -conotoxin peptides or other compounds which have specificity for the $\alpha 3\beta 4$, $\alpha 3\beta 2$ or $\alpha 7$
25 containing subtypes of nAChRs using conventional techniques. One suitable technique involves competitive binding or displacement of the peptide or compound in question with, for example, MII. Peptides or compounds identified in this manner will have the same activity as the compounds used for the screening assay.

-15-

EXAMPLES

The present invention is described by reference to the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below were utilized.

EXAMPLE 1

Chemical Synthesis of α -Conotoxins From *Conus aulicus*

The synthesis of AuIA, AuIB and AuIC conopeptides was separately performed using conventional protection chemistry as described by Cartier et al., 1996a. Briefly, the linear chains were built on Rink amide resin by Fmoc procedures with 2-(1H-benzotriol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate coupling using an ABI model 430A peptide synthesizer with amino acid derivatives purchased from Bachem (Torrence CA). Orthogonal protection was used on cysteines: Cys³ and Cys¹⁶ were protected as the stable Cys(S-acetamidomethyl), while Cys² and Cys⁸ were protected as the acid-labile Cys(S-trityl). After removal of the terminal Fmoc protecting group and cleavage of the peptides from the resins, the released peptides were precipitated by filtering the reaction mixture into -10°C methyl t-butyl ether, which removed the protecting groups except on Cys³ and Cys¹⁶. The peptides were dissolved in 0.1% TFA and 60% acetonitrile and purified by RPLC on a Vydac C₁₈ preparative column (22 x 250 mm) and eluted at a flow rate of 20 mL/min with a gradient of acetonitrile in 0.1% TFA.

The disulfide bridges in the three conopeptides were formed as described in Cartier et al. (1996a). Briefly, the disulfide bridges between Cys² and Cys⁸ were formed by air oxidation which was judged to be complete by analytical RPLC. The monocyclic peptides were purified by RPLC on a Vydac C₁₈ preparative column (22 x 250 mm) and eluted with a gradient of acetonitrile in 0.1% TFA. Removal of S-acetamidomethyl groups and closure of the disulfide bridge between Cys³ and Cys¹⁶ was carried out simultaneously by iodine oxidation. The cyclic peptides were purified by RPLC on a Vydac C₁₈ preparative column (22 x 250 mm) and eluted with a gradient of acetonitrile in 0.1% TFA.

EXAMPLE 2

Biological Activity of α -Conotoxins From *Conus aulicus*

Each of the AuIA, AuIB and AuIC conopeptides were tested for activity on neuronal nAChRs in *Xenopus laevis* oocytes containing different subtypes of nAChRs as described by Cartier

-16-

et al. (1996a). Briefly, oocytes were injected with RNA encoding the various α and β subunits of rat nAChRs and incubated at 25° C for 1-9 days prior to use. Electrophysiological currents were measured using conventional techniques, such as described in Cartier et al. (1996a). Measurements were made for oocytes perfused with acetylcholine as controls and for oocytes incubated with 3 μ M of either AuIA conopeptide, AuIB conopeptide or AuIC conopeptide followed by perfusion with acetylcholine. Each of these conopeptides was active on neuronal nAChRs with a preference for nAChRs of the $\alpha 3\beta 4$ subtype. AuIB was found to be the most potent and selective of the three for the $\alpha 3\beta 4$ subtype. The biological activity of these peptides against the panel of nAChR subtypes is shown in Figure 1.

EXAMPLE 3

Synthesis of Iodinated Tyr-MII

Tyr-MII was prepared in accordance with the procedure of Example 1. Iodination of the Tyr-MII was performed by the Chloramine T method. Briefly, excess peptide was mixed with NaI (either radioactive or nonradioactive version). Chloramine T was then added to initiate the iodination process. Procedure was carried out at a somewhat acidic pH (5.3) to selectively iodinate the Tyr (instead of the His which can also be iodinated at basic pH). Reaction was terminated by the addition of excess ascorbic acid. Mono-iodo and di-iodo Tyr-MII were purified from unmodified peptide using RPLC. The mono- and di-iodo peptides both retain activity as measured by antagonist activity on nAChRs expressed in *Xenopus* oocytes, with a preference for the $\alpha 3\beta 2$ subtype.

EXAMPLE 4

Synthesis and Activity of FAT-MII

FAT-MII was prepared in accordance with the procedure of Example 1 and its activity was measured in accordance with the procedure of Example 2. While MII shows a preference for the $\alpha 3\beta 2$ subtype, FAT-MII shows a preference for the $\alpha 3\beta 4$ subtype.

EXAMPLE 5

Synthesis and Activity of PnIA Analogs

PnIA A10L and PnIA N11S were prepared in accordance with the procedure of Example 1 and their activities were measured in accordance with the procedure of Example 2. The antagonistic

-17-

activity of these peptides as well as peptides PnIA and PnIB for the $\alpha 7$ subtype is shown in Figure 2. Figure 2 shows that PnIA A10L has a higher affinity to the $\alpha 7$ subtype than PnIB which shows preference to this subtype. PnIA shows preference to the $\alpha 3\beta 2$ subtype. The IC_{50} (in nmol) for each peptide with respect to $\alpha 7$ nAChR is as follows: PnIA N11S: 1705; PnIA: 229; PnIB: 61; and PnIA A10L: 12.

EXAMPLE 6

The Effects of α -Conotoxin MII on
Nicotine-stimulated [3 H]-Dopamine Release

The ability of α -conotoxin MII to block nicotine-evoked [3 H]-dopamine release was assessed using rat striatal synaptosomes as described in U.S. Serial No. 08/761,764 or PCT/US97/22350. Three μ M nicotine stimulates the release of [3 H]-dopamine. This release is fully blocked by the non-selective, noncompetitive antagonist mecamylamine. The release is abolished in the absence of external calcium. (-)-Nicotine has previously been shown to increase [3 H]-dopamine release from rat striatal synaptosomes in a concentration-dependent manner with an estimated EC_{50} of 1.6×10^{-7} M (El-Bizri and Clarke, 1994). α -Conotoxin MII blocked 3 μ M nicotine-stimulated [3 H]-dopamine release with a nonsignificant trend toward block at 0.1 nM ($P = 0.08$). At concentrations of 1 nM and above, α -conotoxin significantly blocked nicotine-evoked [3 H]-dopamine release in a dose-dependent manner. Concentrations of α -conotoxin MII of 10 nM and below are expected to be specific for $\alpha 3\beta 2$ receptors (IC_{50} 0.5 nM), whereas concentrations of 100 nM and 1 μ M may have measurable effects on other nAChR subtypes (Cartier et al., 1996a; Cartier et al., 1996b).

EXAMPLE 7

Effects of α -Conotoxin MII on
Depolarization-stimulated [3 H]-Dopamine Release

To further investigate the specificity of effects of α -conotoxin MII, its effects on KCl-induced dopamine release were assessed. One hundred nM α -conotoxin MII, a concentration which blocks 33% of nicotine stimulated dopamine release, had no effect on potassium-stimulated dopamine release.

-18-

EXAMPLE 8

Effects of α -Conotoxin MII on 100 μ M
Nicotine-stimulated [3 H]-Dopamine Release

Previous investigators who have examined the effects of κ -bungarotoxin on nicotine-stimulated dopamine release have reported variable effects. One group reported a 50% inhibition of striatal dopamine release by 100 nM κ -bungarotoxin (Wonnacott et al., 1995). Other investigators have reported complete block of striatal dopamine release by 100 nM κ -bungarotoxin (Grady et al., 1992; Wilkie et al., 1993; Schultz and Zigmond, 1989). One difference between these studies is that the investigators who observed a 50% inhibition used 3 μ M nicotine whereas the investigators who observed complete block used 50 or 100 μ M nicotine. It has been suggested that at 3 μ M, nicotine could be acting on a higher-affinity nicotinic receptor that has low sensitivity to κ -bungarotoxin whereas 50 to 100 μ M nicotine is acting on a lower-affinity nAChR which has a high sensitivity to κ -bungarotoxin (Wonnacott et al., 1995). To more fully compare our results with previously reported results with κ -bungarotoxin, the effects of α -conotoxin MII on 160 nM, 3 μ M and 100 μ M-stimulated dopamine release were tested. One hundred nM α -conotoxin MII blocks 44% of 100 μ M nicotine-stimulated [3 H]-dopamine release compared to 34% of 3 μ M nicotine and 50% of 160 nM nicotine stimulated [3 H]-dopamine release.

EXAMPLE 9

Effects of α -Conotoxin MII on
Nicotine-stimulated Norepinephrine Release

Nicotine evokes the release of [3 H]-norepinephrine in hippocampus. It has previously been reported that nicotine releases striatal dopamine more potently than hippocampal norepinephrine (EC_{50} = 0.16 μ M vs. 6.5 μ M) in synaptosomal preparations (Clarke and Reuben, 1996). Three μ M nicotine was utilized to maximize the chance of seeing an effect by the competitive antagonist α -conotoxin MII. One hundred nM α -conotoxin MII blocked 0% of nicotine-stimulated norepinephrine release which was not statistically different from control. In contrast, 1 μ M α -conotoxin MII blocked 45% (Figure 4). α -Conotoxin MII was also tested on 100 μ M nicotine-stimulated release. One hundred nM MII blocked 11% and 1 μ M blocked 24%.

EXAMPLE 10

Experimental Procedures for
Analysis of Effects of α -Conotoxin AulB

Materials: [^3H]-Dopamine (dihydroxyphenyl-ethylamine, 3,4 [^3H]-) (~ 30 Ci/mmol) and
5 [^3H]-norepinephrine (norepinephrine, levo-[ring-2,5,6- [^3H]-) (~ 42 Ci/mmol) were purchased from
Dupont NEN, Boston, MA. [^3H]-Radioligands were distributed into 5 and 14.1 μCi aliquots
respectively and stored under argon at -80°C . (-)Nicotine hydrogen tartrate was from Sigma.
Pargyline HCl and mecamylamine HCl were from Research Biochemicals International. Prior to
use, all drugs were prepared fresh in superfusion buffer (SB) consisting of 128 mM NaCl, 2.4 mM
10 KCl, 3.2 mM CaCl_2 , 1.2 mM KH_2PO_4 , 0.6 mM MgSO_4 , 25 mM HEPES, 10 mM D-glucose, 1 mM
L-ascorbic acid, 0.1 mM pargyline, 0.1 mM BSA with the pH adjusted to 7.5 with NaOH.

Animals: Male Sprague-Dawley rats, weighing 200-400 g, were maintained on a 12/12h
light/dark cycle. Rats were drug-naïve and housed three per cage, and food and water were available
ad libitum.

15 Synaptosomal Preparation and [^3H]-Radioligand Preloading: Synaptosomes were prepared
as previously described (Kulak et al., 1997). A crude P2 synaptosomal fraction was resuspended
in SB (0.5 ml/100 mg wet tissue weight) containing 0.12 μM [^3H]-dopamine for striatal tissue or
0.2 μM [^3H]-norepinephrine for hippocampal tissue and incubated at 37°C for 10 min. The loaded
synaptosomes were centrifuged at 1000g for 5 min at room temperature (24°C), and the pellet was
20 gently resuspended in 2.0 ml of SB. The high [K^+]-stimulated release solution was SB in which the
[K^+] was elevated to 22.4 mM and the [Na^+] was decreased to 108 mM.

Superfusion: The assay system was as previously described (Kulak et al., 1997). Briefly,
the system had twelve identical channels connected to a pump which continuously pulled the
superfusate through individual filter holders containing the synaptosomes at a rate of 0.5 ml/min.
25 Teflon® TFE tubing and Teflon®-coated parts were used upstream of the synaptosomes to avoid
plasticizers such as Tinuvin 770 (a common light and UV radiation stabilizer used in a wide range
of plastics) known to block neuronal nAChRs (Papke et al., 1994).

Following a preliminary superfusion period of 13 minutes for assays containing α -conotoxin
AulB or 31 minutes for all other toxins, a 1 min (0.5 ml) pulse of synaptosomal buffer with or
30 without agonist was delivered simultaneously to all channels by switching on solenoids. Nicotine
concentration was 3 μM in dopamine-release experiments and 100 μM in norepinephrine-release
experiments. Two-minute fractions per channel were collected in polypropylene minivials

-20-

containing 4.0 ml of scintillation fluid. Following the collection period, the filters holding the synaptosomes were removed to determine the residual radioactivity. A liquid scintillation counter (Beckman LS9800, 57.2% efficiency) was used to determine tritium levels.

Data Analysis: It has previously been shown that tritium released by nAChR agonists or b
5 depolarizing concentrations of KCl is directly proportional to total radioligand released (Rapier et al., 1988). Thus, levels of tritium released is assumed to correspond directly to amount of radioligand released.

Release is calculated as: (dpm in the peak fractions - baseline release)/baseline release. Baseline release is defined as the average of two pre- and two post-release fractions. Release is
10 normalized as a percentage of total-agonist-stimulated release. Agonist-stimulated release with superfusate containing different α -conotoxin concentrations were compared to those of controls without toxin and analyzed for statistically significant mean differences using a t-test on raw (non-normalized) data with SPSS software (SPSS, Chicago, Illinois).

EXAMPLE 11

Effects of α -Conotoxin AulB on Nicotine-stimulated Norepinephrine Release

Presynaptic nicotinic receptors are known to be involved in the release of various neurotransmitters including norepinephrine and dopamine. The effects of α -conotoxin AulB and
20 other α -conotoxins were assessed in this regard. Nicotine-stimulated norepinephrine or dopamine release was analyzed using synaptosomes from rat hippocampus or from rat striatum, respectively. AulB at 1 and 5 μ M block a portion of nicotine-stimulated norepinephrine release but not that of dopamine release. The converse result is obtained using the $\alpha 3\beta 2$ selective α -conotoxin MII. MII blocks nicotine-stimulated dopamine release but has no effect on nicotine-stimulated norepinephrine
25 release. In addition, it was found that the $\alpha 7$ selective α -conotoxin ImI and $\alpha 1$ selective MI all fail to block nicotine-stimulated norepinephrine release. Similarly, it was found that ImI and MI also fail to block nicotine-stimulated release of dopamine. These results are shown in Figures 3A and 3B. The significance of this data is as follows: (a) norepinephrine release: 1 μ M AulB, $p = 0.001$; 5 μ M AulB, $p < 0.001$; MII, $p = 0.75$; ImI, $p = 0.64$; MI, $p = 0.5$; (b) dopamine release: 1 μ M
30 AulB, $p = 0.93$; 5 μ M AulB, $p = 0.68$; MII, $p < 0.001$; ImI, $p = 0.24$; MI, $p = 0.85$.

The specificity of α -conotoxin AulB's block of norepinephrine release was further assessed by testing its effects on high $[K^+]$ -induced norepinephrine release. Hippocampal synaptosomes were

-21-

loaded with [³H]-norepinephrine and pre-incubated with or without α -conotoxin AuIB. Synaptosomes were subsequently depolarized with a one minute pulse of synaptosomal buffer that contained high K⁺ with or without AuIB. Concentrations of AuIB which significantly block nicotine-stimulated norepinephrine release had no effect on depolarization-stimulated norepinephrine release. One μ M AuIB responses = $92.2 \pm 3.7\%$, $p = 0.5$; 5 μ M AuIB response = $99.0 \pm 4.6\%$, $p = 0.95$.

EXAMPLE 12

Inhibition of SCLC Proliferation by α -Conotoxins

Small cell lung carcinoma (SCLC) cells have been found to express cholinergic nicotinic receptors (Maneckjee et al., 1990; Chini et al., 1992; Tarroni et al., 1992; Schuller et al., 1990). These SCLC nicotinic receptors have been shown to be of neuronal type (Chini et al., 1992; Tarroni et al., 1992). Nicotine and cytosine each stimulate the release of 5-hydroxytryptamine (5HT or serotonin) which acts as a potent mitogen in SCLC cells (Maneckjee et al., 1990; Cattaneo et al., 1993). α -Conotoxin MI has been found to block the nicotine or cytosine induced release of serotonin and at a concentration of 1 μ M it completely antagonized the nicotine and cytosine stimulation of SCLC proliferation (Codignola et al., 1994). α -Conotoxins which bind to neuronal type nicotinic receptors are suitable for preventing the proliferation of tumors such as SCLC and can be used therapeutically to inhibit such proliferation as described below. These α -conotoxins can also be used diagnostically for detecting the presence and/or location of small-cell lung tumors as described below. Although Codignola et al. (1994) report that α -conotoxin MI binds to these SCLC receptors, α -conotoxin MI is not suitable for therapeutic or diagnostic use since it also binds to neuromuscular receptors and can cause paralysis which could lead to death. α -Conotoxins which do not bind to neuromuscular receptors or which have a much lower affinity for such receptors as compared to the nicotinic neuronal receptors are suitable for therapeutic or diagnostic purposes. Such peptides include the α -conotoxins MII and Iml.

EXAMPLE 13

Diagnosis of SCLC Using α -Conotoxins

α -Conotoxins which bind to SCLC nicotinic receptors can be used for diagnosing SCLC tumors in patients. Suitable α -conotoxins include MII, Iml, PnIB and PnIA A10L. Administration of a labeled conotoxin to a patient will reveal the presence of SCLC cells if any are present. The

-22-

α -conotoxin is labeled with a radioactive marker, preferably iodine, e.g., ^{131}I or ^{125}I . Labeling can be performed by standard techniques well known in the art. Alternatively, a Tyr residue can be added to the N-terminus and iodinated as described above. The labeled toxin is administered intravenously in a range of 5-50 nmoles, preferably about 25 nmoles. The label is then detected by standard techniques well known in the art. The labeled toxins will bind to SCLC cells and also may bind to autonomic ganglia. However, the locations of autonomic ganglia are known and can be distinguished from signals resulting from binding of the labeled toxin to SCLC cells.

EXAMPLE 14

Therapeutic Use of α -Conotoxins to Treat SCLC Tumors

α -Conotoxins which bind to SCLC nicotinic receptors can be used therapeutically to treat patients with SCLC tumors. Suitable conotoxins are those which do not bind strongly to muscle receptors, e.g., MII, ImI, PnIB and PnIA A10L. Patients who have been diagnosed with SCLC can have a suitable conotoxin administered, preferably intravenously or intramuscularly. A dose of 200-2000 nanomoles, preferably about 500 nanomoles, is administered. The dosing schedule depends on the in vivo stability of the specific conotoxin used. In general conotoxins are relatively resistant to degradation and may last on the order of a few days. Therefore a typical dosing schedule may be anywhere from twice per day to once every few days, this being dependent on the biological lifetime of the specific conotoxin used.

It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will be apparent to the artisan that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

LIST OF REFERENCES

- Bodansky et al. (1966). *Chem. Ind.* 38:1597-98.
Cartier, G.E. et al. (1996a). *J. Biol. Chem.* 271:7522-7528.
Cartier, G.E. et al. (1996b). *Soc. Neurosci. Abst.* 22:268.
Cattaneo, M.G. et al. (1993). *Cancer Res.* 53: 566-5568.
Chini, B. et al. (1992). *Proc. Natl. Acad. Sci. USA* 89:1572-1576.

- Clarke, P.B.S. and Reuben, M. (1996). *Br. J. Pharmacol.* 111:695-702.
- Codignola, A. et al. (1994). *FEBS Lett.* 342:286-290.
- Cruz, L.J. et al. (1976). *Verliger* 18:302-308.
- 5 Cruz, L.J. et al. (1987). *Conus geographus* toxins that discriminate between neuronal and muscle sodium channels. *J. Biol. Chem.* 260:9280-9288.
- El-Bizri, H. and Clarke, P.B.S. (1994). *Br. J. Pharmacol.* 111:406-413.
- Grady, S. et al. (1992). *J. Neurochem.* 59:848-856.
- Gray, R. et al. (1996). *J. Neurochem.* 59:848-856.
- 10 Haack, J.A. et al. (1990). Contryphan-T: a gamma-carboxyglutamate containing peptide with N-methyl-d-aspartate antagonist activity. *J. Biol. Chem.* 265:6025-6029.
- Horiki, K. et al. (1978). *Chemistry Letters* 165-68.
- Kahn, R.S. and Davis, K.L. (1995). New Developments in Dopamine and Schizophrenia. In *Psychopharmacology: The Fourth Generation of Progress*, Bloom, F.E. and Kupfer, D.J. (eds), Raven Press, New York, pp 1193-1203.
- 15 Kaiser et al. (1970). *Anal. Biochem.* 34:595.
- Kapoor (1970). *J. Pharm. Sci.* 59:1-27.
- Kornreich, W.D. et al. (1986). U.S. Patent No. 4,569,967.
- Kulak, J.M. et al. (1997). α -Conotoxin MII blocks nicotine-stimulated dopamine release in rat striatal synaptosomes. *J. Neurosci.* 17:5263-5270.
- 20 Lapchak, P.A. et al. (1989). *J. Neurochem.* 52:483-491.
- Maneckjee, R. and Minna, J.D. (1990). *Proc. Natl. Acad. Sci. USA* 87:3294-3298.
- Martinez, J.S. et al. (1995). *Biochem.* 34:14519-14526.
- McGehee, D.S. and Role, L.W. (1995). *Annu. Rev. Physiol.* 57:521-546.
- 25 Mena, E.E. et al. (1990). Contryphan-G: a novel peptide antagonist to the N-methyl-D-aspartic acid (NMDA) receptor. *Neurosci. Lett.* 118:241-244.
- Methoden der Organischen Chemie (Houben-Weyl): Synthese von Peptiden*, E. Wunsch (Ed.), Georg Thieme Verlag, Stuttgart, Ger. (1974).
- Mongeau, R. et al. (1997). The serotonergic and noradrenergic systems of the hippocampus: their interactions and the effects of antidepressant treatments. *Brain Res. Rev.* 23:145-195.
- 30 Mulle, C. et al. (1991). *J. Neurosci.* 11:2588-2597.
- Nishiuchi, Y. et al. (1993). Synthesis of gamma-carboxyglutamic acid-containing peptides by the Boc strategy. *Int. J. Pept. Protein Res.* 42:533-538.
- Olivera, B.M. et al. (1984). U.S. Patent 4,447,356.
- 35 Olivera, B.M. et al. (1985). Peptide neurotoxins from fish-hunting cone snails. *Science* 230:1338-1343.

- Papke, R.L. et al. (1994). Inhibition of nicotinic acetylcholine receptors by bis(2,2,6,6-tetramethyl-4r-piperidinyl)sebacate (Tinuvin 770), an additive to medical plastics. *J. Pharmacol. Exp. Ther.* 268:718-726.
- 5 Pontieri, F.E. et al. (1996). Effects of nicotine on the nucleus accumbens and similarity to those of addictive drugs. *Nature* 383:255-257.
- Rapier, C. et al. (1988). *J. Neurochem.* 50:1123-1130.
- Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, PA (1990).
- Rivier, J.E. et al. (1978). *Biopolymers* 17:1927-38.
- 10 Rivier, J.E. et al. (1987). Total synthesis and further characterization of the gamma-carboxyglutamate-containing 'sleeper' peptide from *Conus geographus*. *Biochem.* 26:8508-8512.
- Roberts et al. (1983). *The Peptides* 5:342-429.
- Rowell, P.P. and Winkler, D.L. (1984). *J. Neurochem.* 43:1593-1598.
- Sacaan, A.I. et al. (1995). *J. Pharmacol. Exp. Therapeutics* 274:224-230.
- 15 Sambrook, J. et al. (1979). *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Schatzberg, A.F. and Nemeroff, C.B (eds) (1995). "Section II: Classes of psychiatric drugs: animal and human pharmacology." In *Textbook of Pharmacology*, Meltzer, H.Y. (ed), American Psychiatric Press, Inc., Washington, D.C., pp. 141-438.
- 20 Schroder & Lubke (1965). *The Peptides* 1:72-75, Academic Press, NY.
- Schuller, H.M. et al. (1990). *Life Sci.* 47:571-578.
- Schultz, D.W. and Zigmond, R.E. (1989). *Neurosci. Lett.* 98:310-316.
- Shon, K.-J. et al. (1994). *Biochemistry* 33:11420-11425.
- Sine, S.M. and Claudio, T. (1991). *J. Biol. Chem.* 266:19369-19377.
- 25 Stewart and Young, *Solid-Phase Peptide Synthesis*, Freeman & Co., San Francisco, CA (1969).
- Tarroni, P. et al. (1992). *FEBS Lett.* 312:66-70.
- Vale et al. (1978). U.S. Patent 4,105,603.
- Vidal, C. and Changeux, J.-P. (1993). *Neuroscience* 56:23-32.
- Wilkie, G.I. et al. (1993). *Biochem. Soc. Trans.* 21:429-431.
- 30 Wonnacott, S. et al. (1995). "Presynaptic nicotinic autoreceptors and heteroreceptors in the CNS". In *Effects of Nicotine on Pharmacological Systems II*, P.B.S. Clarke et al. (eds.), Birkhäuser Verlag, Basel, pp. 87-94.
- Zhou L.M., et al. (1996a). Synthetic Analogues of Contryphan-G: NMDA Antagonists Acting Through a Novel Polyamine-Coupled Site. *J. Neurochem.* 66:620-628.
- 35 U.S. Patent 3,972,859 (1976).

U.S. Patent 3,842,067 (1974).

U.S. Patent 3,862,925 (1975).

PCT Published Application WO 96/11698

WHAT IS CLAIMED IS:

1. Use of an α -conotoxin peptide for preparing a pharmaceutical composition for treating disorders regulated at neuronal nicotinic acetylcholine receptors (nAChRs), said α -conotoxin peptide having the general formula
5
Xaa₁-Xaa₂-Cys-Cys-Xaa₃-Xaa₄-Pro-Xaa₅-Cys-Xaa₆-Cys (SEQ ID NO:1)
wherein Xaa₁ is des-Xaa₁, Tyr, mono-iodo-Tyr or di-iodo-Tyr, Xaa₂ is any amino acid, Xaa₃ is any amino acid, Xaa₄ is any amino acid, Xaa₅ is any amino acid and Xaa₆ represents a peptide of 3-7 amino acids or a pharmaceutically acceptable salt thereof, with the proviso
10 that when the disorder is small cell lung carcinoma, then the α -conotoxin peptide is not a peptide having an amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:13.
2. The use of claim 1 wherein Xaa₁ is Tyr, mono-iodo-Tyr or di-iodo-Tyr.
- 15 3. The use of claim 1 or 2 wherein said nAChR is an $\alpha 3 \beta 2$ -containing nAChR.
4. The use of claim 1 or 2 wherein said nAChR is an $\alpha 3 \beta 4$ -containing nAChR.
5. The use of claim 1 or 2 wherein said nAChR is an $\alpha 7$ -containing nAChR.
- 20 6. The use of any one of claims 1 to 5 wherein said disorder is a cardiovascular disorder.
7. The use of any one of claims 1 to 5 wherein said disorder is a gastric motility disorder.
- 25 8. The use any one of claims 1 to 5 wherein said disorder is urinary incontinence.
9. The use of any one of claims 1 to 5 wherein said disorder is nicotine addiction.
10. The use of any one of claims 1 to 5 wherein said disorder is a mood disorder.
- 30 11. The use of any one of claims 1 to 5 wherein said disorder is small cell lung carcinoma.

-27-

12. A substantially pure α -conotoxin peptide selected from the group consisting of:
Gly-Cys-Cys-Ser-Leu-Pro-Pro-Cys-Ala-Leu-Asn-Asn-Pro-Asp-Tyr-Cys (SEQ ID NO:10);
Gly-Cys-Cys-Ser-Leu-Pro-Pro-Cys-Ala-Ala-Ser-Asn-Pro-Asp-Tyr-Cys (SEQ ID NO:11);
Tyr-Gly-Cys-Cys-Ser-Asn-Pro-Val-Cys-His-Leu-Glu-His-Ser-Asn-Leu-Cys (SEQ ID
5 NO:3); and
Gly-Cys-Cys-Ser-Asn-Pro-Val-Cys-Phe-Ala-Thr-His-Ser-Asn-Leu-Cys (SEQ ID NO:4).
13. The α -conotoxin peptide of claim 12 having the formula Gly-Cys-Cys-Ser-Leu-Pro-Pro-
Cys-Ala-Leu-Asn-Asn-Pro-Asp-Tyr-Cys (SEQ ID NO:10).
10
14. The α -conotoxin peptide of claim 12 having the formula Gly-Cys-Cys-Ser-Leu-Pro-Pro-
Cys-Ala-Ala-Ser-Asn-Pro-Asp-Tyr-Cys (SEQ ID NO:11).
15. The α -conotoxin peptide of claim 12 having the formula Tyr-Gly-Cys-Cys-Ser-Asn-Pro-
Val-Cys-His-Leu-Glu-His-Ser-Asn-Leu-Cys (SEQ ID NO:3).
15
16. The α -conotoxin peptide of claim 12 having the formula Gly-Cys-Cys-Ser-Asn-Pro-Val-
Cys-Phe-Ala-Thr-His-Ser-Asn-Leu-Cys (SEQ ID NO:4).
- 20 17. The α -conotoxin peptide of claim 13 wherein at least one of the Pro residues is replaced with
hydroxyproline.
18. The α -conotoxin peptide of claim 13 or 14 wherein a Tyr residue is incorporated on the N-
terminus.
25
19. The α -conotoxin peptide of claim 18 wherein the Tyr residue is substituted with one or two
iodines.
20. The α -conotoxin peptide of claim 14 wherein at least one of the Pro residues is replaced with
hydroxyproline.
30

-28-

21. The α -conotoxin peptide of claim 14 or 20 wherein a Tyr residue is incorporated on the N-terminus.
22. The α -conotoxin peptide of claim 21 wherein the Tyr residue is substituted with one or two iodines.
23. The α -conotoxin peptide of claim 15 wherein the Pro residue is replaced with hydroxyproline.
24. The α -conotoxin peptide of claim 15 or 23 wherein the Tyr residue is substituted with one or two iodines.
25. The α -conotoxin peptide of claim 16 wherein the Pro residue is replaced with hydroxyproline.
26. The α -conotoxin peptide of claim 16 or 25 wherein a Tyr residue is incorporated on the N-terminus.
27. The α -conotoxin peptide of claim 26 wherein the Tyr residue is substituted with one or two iodines.
28. A substantially pure α -conotoxin peptide having the formula Xaa-peptide, wherein Xaa is Tyr, mono-iodo-Tyr or di-iodo-Tyr and peptide is selected from the group consisting of (a) a peptide having the amino acid sequence set forth in SEQ ID NO:5, (b) a peptide having the amino acid sequence set forth in SEQ ID NO:7, (c) a peptide having the amino acid sequence set forth in SEQ ID NO:8, (d) a peptide having the amino acid sequence set forth in SEQ ID NO:9, (e) a peptide having the amino acid sequence set forth in SEQ ID NO:12 and (f) a peptide having the amino acid sequence set forth in SEQ ID NO:13.
29. The α -conotoxin peptide of claim 28 wherein at least one of the Pro residues in the peptide is replaced with hydroxyproline.

-29-

30. The α -conotoxin peptide of claim 28 or 29 wherein a Trp residue in the peptide is replaced with bromotryptophan.




FIG. 1

SUBSTITUTE SHEET (RULE 26)




FIG. 2

SUBSTITUTE SHEET (RULE 26)




FIG. 3A




FIG. 3B

SUBSTITUTE SHEET (RULE 26)

-1-

SEQUENCE LISTING

5 <110> Olivera, Baldomero M.
 McIntosh, J. Michael
 Yoshikami, Doju
 Cartier, G. Edward
 Luo, Siqin
 University of Utah Research Foundation

10 <120> Uses of Alpha-Conotoxin Peptides
 <130> Uses of Alpha-Conotoxins

15 <140>
 <141>

<150> US 60/080,588
 <151> 1998-04-03

20 <150> US 60/070,153
 <151> 1997-12-31

<160> 13

25 <170> PatentIn Ver. 2.0
 <210> 1
 <211> 11
 <212> PRT

30 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:generic
 alpha-conotoxin sequence

35 <220>
 <221> PEPTIDE
 <222> (1)..(6)
 <223> Xaa at residue 1 is des-Xaa, Tyr, mono-iodo-Tyr or
 40 di-iodo-Tyr; Xaa at residue 2 is any amino acid;
 Xaa at residue 5 is any amino acid; Xaa at residue
 6 is any amino acid.

<220>
 45 <221> PEPTIDE
 <222> (8)..(10)
 <223> Xaa at residue 8 may be any amino acid; Xaa at
 residue 10 is a peptide having 3-7 amino acids.

50 <400> 1
 Xaa Xaa Cys Cys Xaa Xaa Pro Xaa Cys Xaa Cys
 1 5 10

55 <210> 2
 <211> 16
 <212> PRT
 <213> Conus magus

60 <400> 2
 Gly Cys Cys Ser Asn Pro Val Cys His Leu Glu His Ser Asn Leu Cys
 1 5 10 15

-2-

```

5      <210> 3
      <211> 17
      <212> PRT
      <213> Artificial Sequence

      <220>
      <223> Description of Artificial Sequence:Tyr derivative
            of C. magus MII

10     <400> 3
      Tyr Gly Cys Cys Ser Asn Pro Val Cys His Leu Glu His Ser Asn Leu
            1             5             10             15
      Cys

15

      <210> 4
      <211> 16
      <212> PRT
      <213> Artificial Sequence

      <220>
      <223> Description of Artificial Sequence:FAT derivative
            of C. magus MII

25     <400> 4
      Gly Cys Cys Ser Asn Pro Val Cys Phe Ala Thr His Ser Asn Leu Cys
            1             5             10             15

30

      <210> 5
      <211> 16
      <212> PRT
      <213> Conus aulicus

35     <400> 5
      Gly Cys Cys Ser Tyr Pro Pro Cys Phe Ala Thr Asn Ser Asp Tyr Cys
            1             5             10             15

40

      <210> 6
      <211> 17
      <212> PRT
      <213> Artificial Sequence

45     <220>
      <223> Description of Artificial Sequence:Tyr derivative
            of C. aulicus AuIA

50     <400> 6
      Tyr Gly Cys Cys Ser Tyr Pro Pro Cys Phe Ala Thr Asn Ser Asp Tyr
            1             5             10             15

55     Cys

60     <210> 7
      <211> 15
      <212> PRT
      <213> Conus aulicus

```

<400> 7

5

<210> 8

<211> 16

<212> PRT

<213> Conus aulicus

10

<400> 8

Gly Cys Cys Ser Tyr Pro Pro Cys Phe Ala Thr Asn Ser Gly Tyr Cys
1 5 10 15

15

<210> 9

<211> 16

<212> PRT

<213> *Conus purpurascens*

20

<400> 9

Gly Cys Cys Ser Leu Pro Pro Cys Ala Ala Asn Asn Pro Asp Tyr Cys
1 5 10 15

25

<210> 10

<211> 16

<212> PR'T

<213> Artificial Sequence

30

$\langle 220 \rangle$

<223> Description of Artificial Sequence: A10L derivative
of *C. purpurascens* PnIA

35

<400> 10

Gly Cys Cys Ser Leu Pro Pro Cys Ala Leu Asn Asn Pro Asp Tyr Cys
1 5 10 15

40

<210> 11

<211> 16

<212> PRT.

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence:N11S derivative
of *C. purpurascens* PnIA

<400> 11

Gly Cys Cys Ser Leu Pro Pro Cys Ala Ala Ser Asn Pro Asp Tyr Cys
1 5 10 15

55

<210> 12

<211> 16

<212> PRT

<213> *Conus purpurascens*

60

<400> 12

Gly Cys Cys Ser Leu Pro Pro Cys Ala Leu Ser Asn Pro Asp Tyr Cys
1 5 10 15

-4-

<210> 13
<211> 12
<212> PRT
<213> Conus imperialis

5

<400> 13
Gly Cys Cys Ser Asp Pro Arg Cys Ala Trp Arg Cys
1 5 10

10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/27367

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : A61K 38/00, 38/04 US CL : 514/9, 13, 14; 530/317, 326, 327 According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/9, 13, 14; 530/317, 326, 327 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, CAS REGISTRY sequence search.														
C. DOCUMENTS CONSIDERED TO BE RELEVANT														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	KULAK, J.M. Alpha-Conotoxin MII Blocks Nicotine-Stimulated Dopamine Release From Rat Striatal Synaptosomes J. Neurosci. 15 July 1997, Vol. 17, No. 14, page 5263-5270, especially pages 5263 and 5266.	1,28												
X	FAINZILBER, M. New Mollusc-Specific Alpha-Conotoxins Block Aplysia Neuronal Acetylcholine Receptors. Biochemistry. 1994, Vol. 33, No. 32, pages 9523-9529, especially page 9523, Abstract, line 7.	28												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"><tr><td>* Special categories of cited documents:</td><td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td></tr><tr><td>*A* document defining the general state of the art which is not considered to be of particular relevance</td><td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td></tr><tr><td>*E* earlier document published on or after the international filing date</td><td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td></tr><tr><td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td><td>*A* document member of the same patent family</td></tr><tr><td>*O* document referring to an oral disclosure, use, exhibition or other means</td><td></td></tr><tr><td>*P* document published prior to the international filing date but later than the priority date claimed</td><td></td></tr></table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family													
O document referring to an oral disclosure, use, exhibition or other means														
P document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 15 APRIL 1999		Date of mailing of the international search report 12 MAY 1999												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer FABIAN JAMEISON Telephone No. (703) 305-0509												

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/27367

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TAVAZOIE, S.F. et al. Differential Block of Nicotinic Synapses on B Versus C Neurones in Sympathetic Ganglia of Frog by alpha-Conotoxins MII and ImI British J. Pharmacol. 1997, Vol. 120, No. 6, page 996-1000, especially page 996, lines 5-6.	28
X	US 5,514,774 A (OLIVERA et al) 07 May 1996, col. 61, lines 31-34, and column 29.	12, 15 and 28
X	US 5,633,347 A (OLIVERA et al) 27 May 1997, col. 6, lines 10-12.	12 and 15
X	WO 95/11256 A (UNIVERSITY OF UTAH RESEARCH FOUNDATION) 27 April 1995, page 43.	1, 12, 18, 26 and 28
A	US 5,432,155 A (OLIVERA et al) 11 June 1995, entire article.	12-28

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU 99/00843

A. CLASSIFICATION OF SUBJECT MATTER		
Int Cl ⁶ : C07K 7/08; C07H 21/02, 21/04; A61K 31/195		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) ANGIS. Homology search on seq. ID. No. 1; STN File DGene. Seq. ID. No. 1; STN Files CA, Medline, WPIDS. Keywords Conotoxin and Conus (W) tulipa		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A, 9824462 (University of Utah Research Foundation), 11 June 1998. See whole document.	1-23
X	WO, A, 9511256 (University of Utah Research Foundation), 27 April 1995. See whole document.	1-23
P.X	WO, A, 9921878 (University of Utah Research Foundation et. al.), 6 May 1999. See whole document.	1-23
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents: "A" Document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "(C)" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 28 October 1999		Date of mailing of the international search report 11 NOV 1999
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200 WODEN ACT 2606 AUSTRALIA mail address: pct@jpaaustralia.gov.au No.: (02) 6285 3929		Authorized officer L.F. MCCAFFERY Telephone No.: (02) 6283 2573

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU 99/00843

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO, A, 9933482 (University of Utah Research Foundation), 8 July 1999. See whole document.	1-23

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/AU 99/00843

Box I Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1, 4-23
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claim 1 and appended claims define p-conotoxin derivatives. This designation appears an arbitrary form of nomenclature without any established meaning in the field. Accordingly the search has been based substantially on the sequence of claim 2.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box II Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Information on patent family members

International application No.
PCT/AU 99/00843

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	98/24462	AU	56909/98	US	5780433	US	5922679
		US	5929034				
WO	95/11256	AU	10831/95	CA	2172989	EP	728146
		US	5514774	US	5589340	US	5595972
WO	99/21878	AU	11143/99				

END OF